

INPLASY

Effects of delirium in hospitalized children: a systematic review and meta-analysis

INPLASY202370011

doi: 10.37766/inplasy2023.7.0011

Received: 03 July 2023

Published: 03 July 2023

Corresponding author:

Feng Xianqiong

fengxianqiong66@126.com

Author Affiliation:

West China School of Nursing,
Sichuan University/Department Of
Pediatric Intensive Care Unit, West
China Second University Hospital,
Sichuan University.

Yang, C¹; Feng, XQ².

ADMINISTRATIVE INFORMATION

Support - There is no source of financial support.

Review Stage at time of this submission - Formal screening of search results against eligibility criteria.

Conflicts of interest - None declared.

INPLASY registration number: INPLASY202370011

Amendments - This protocol was registered with the International Platform of Registered Systematic Review and Meta-Analysis Protocols (INPLASY) on 03 July 2023 and was last updated on 03 July 2023.

INTRODUCTION

Review question / Objective To explore the influence of delirium on hospitalization outcome in children and whether it affects the quality of life. P: Children; I: Delirium occurred during hospitalization; C: No delirium occurred during hospitalization; O: Mortality, Length of ICU Stay, Mechanical Ventilation rate in hospital and Quality of life; S: Prospective or Cohort study.

Condition being studied Delirium is an acute cognitive disorder, and it has been confirmed that the occurrence of delirium can bring adverse effects on adults, such as prolonged hospital stay, increased mortality, cognitive decline, etc., but whether the same results in children are unclear, which requires further research.

METHODS

Participant or population Hospitalized children.

Intervention Children with delirium during hospitalization.

Comparator Children without delirium during hospitalization.

Study designs to be included Prospective or Cohort study.

Eligibility criteria None.

Information sources Pubmed, embase, cochrane, CINAHL.

Main outcome(s) Mortality, Length of ICU Stay, Mechanical Ventilation rate in hospital and Quality of life.

Quality assessment / Risk of bias analysis the Newcastle-Ottawa Scale, NOS.

Strategy of data synthesis Baseline data were presented using mean \pm standard deviation or median (IQR), and heterogeneity was assessed using $P + I^2$, if heterogeneity was small, using a fixed-effects model, and heterogeneity was greater using a random-effects model to pool the studies, publication offset was assessed by drawing funnel plots, and combined effect sizes were performed with STATA version 17 and RevMan version.

Subgroup analysis Subgroup analysis of outcome according to mortality, mechanical ventilation rate, length of hospital stay and quality of life.

Sensitivity analysis Stata software was used to Sensitivity analysis, by deleting one of the articles after the effect size changes to reflect the sensitive situation.

Country(ies) involved China.

Keywords delirium; children; mortality; mechanical ventilation; length of hospital stay; quality of life.

Contributions of each author

Author 1 - Yang Cheng.

Email: 137864365@qq.com

Author 2 - Feng Xianqiong.

Email: fengxianqiong66@126.com