

# INPLASY

## Association between air pollution and frailty: a systematic review and meta-analysis

INPLASY202410101

doi: 10.37766/inplasy2024.1.0101

Received: 24 January 2024

Published: 24 January 2024

Ding, QL<sup>1</sup>; Feng, Y<sup>2</sup>; Sun, ZY<sup>3</sup>; Kou, CG<sup>4</sup>; Bai, W<sup>5</sup>.

### Corresponding author:

Changgui Kou

koucg@jlu.edu.cn

### Author Affiliation:

Jilin University.

### ADMINISTRATIVE INFORMATION

**Support** - None.

**Review Stage at time of this submission** - Formal screening of search results against eligibility criteria.

**Conflicts of interest** - None declared.

**INPLASY registration number:** INPLASY202410101

**Amendments** - This protocol was registered with the International Platform of Registered Systematic Review and Meta-Analysis Protocols (INPLASY) on 24 January 2024 and was last updated on 24 January 2024.

## INTRODUCTION

**Review question / Objective** What is the relationship between air pollution and frailty?

**Condition being studied** Frailty, a geriatric syndrome characterized by decreased physiological reserves and increased vulnerability, is emerging as one of the most studied clinical conditions in geriatrics due to its reversibility. Air pollution is known as a risk factor for several diseases typically in older adults, such as respiratory diseases and cardiovascular diseases, which are highly associated with frailty. Given this background, our purpose is to summarize the relationship between air pollution and frailty.

## METHODS

**Search strategy** We have searched the following electronic bibliographic database from inception to Nov 15, 2023: PubMed, Web of Science, Embase, and Scopus. The primary search keywords were based on “air pollution” and “frailty”. In “air pollution”, we included the terms “air pollution”, “air pollutants”, “contaminated air” and so on. Keywords in “frailty” were “frail\*” and “frail elderly”. There will be no language restrictions. After removing duplicates, titles, abstracts, and full texts will be screened for eligibility by two authors. Any disagreement will be resolved by consensus with a third author or by discussion with the corresponding author.

---

**Participant or population** Studies conducted in human population without geographical restrictions.

**Intervention** Studies that focused on participants exposure to air pollutants.

**Comparator** Studies comparing individuals exposed to different levels of air pollutants.

**Study designs to be included** We will include both cohort studies and cross-sectional studies.

**Eligibility criteria** Exclusion criteria included irrelevant studies, review, conference, letter, book chapter, editorial material and studies whose effect sizes were not reported.

**Information sources** PubMed, Web of Science, Embase, and Scopus.

**Main outcome(s)** The association between exposure to air pollution and frailty risk (i.e., OR, RR, or HR).

**Quality assessment / Risk of bias analysis** Cohort study quality was assessed using Newcastle-Ottawa Scale (NOS) with overall NOS $\geq$ 7 rating as high-quality. Cross-sectional study quality was assessed using an eight-item assessment instrument for epidemiological studies. Instrument criteria included the following: 1) Target population is clearly defined; 2) Probability or entire population sampling is used; 3) Response rate is >80%; 4) Non-responders are clearly described; 5) Sample is representative of the target population; 6) Data collection methods are standardized; 7) Validated criteria are used to assess the presence of frailty; 8) Estimates of prevalence are given with confidence intervals and detailed by subgroup (if applicable). Total scores ranged from 0 to 8 points. Studies with a total of 7–8 points were considered as ‘high quality’ while those with 4–6 points were of ‘moderate quality’ and those with 0–3 points were ‘low quality’.

**Strategy of data synthesis** Data analyses will be conducted with R; “meta” and “forestplot” modules will be used. Prevalence data will be converted using a Freeman–Tukey double-arcsine transformation. OR and 95% CI of frailty associated with exposure to target air pollutants will be calculated using fixed-effects model or random-effects model. The I<sup>2</sup> statistic was used to evaluate study heterogeneity; values above 50% indicated high heterogeneity.

**Subgroup analysis** Subgroup analysis will be conducted based on categorical variables, such as geographic region, country income level, study design, male ratio, sample size, or other variables, if applicable.

**Sensitivity analysis** Sensitivity analysis will be performed by examining the reliability of results after excluding each study one by one, if applicable.

**Country(ies) involved** China.

**Keywords** Air pollution, frailty, meta-analysis.

#### **Contributions of each author**

Author 1 - Qianlu Ding.

Email: dingfl23@mails.jlu.edu.cn

Author 2 - Yuan Feng.

Email: fengy23@mails.jlu.edu.cn

Author 3 - Zhouyang Sun.

Email: zysun23@mails.jlu.edu.cn

Author 4 - Changgui Kou.

Email: koucg@jlu.edu.cn

Author 5 - Wei Bai.

Email: baiwei@jlu.edu.cn