

INPLASY

Risk of Flare on Drug Therapy in Moderate-to-Severe Atopic Dermatitis: A Network Meta-Analysis

INPLASY202460098

doi: 10.37766/inplasy2024.6.0098

Received: 26 June 2024

Published: 26 June 2024

Jaalouk, D; Humeda, J; Pulumati, A; Algarin, YA; Nouri, K.

Corresponding author:

Dana Jaalouk

danajaalouk@hotmail.com

Author Affiliation:

Department of Dermatology and Cutaneous Surgery, University of Miami Leonard M. Miller School of Medicine, Miami, FL.

ADMINISTRATIVE INFORMATION

Support - None.

Review Stage at time of this submission - Preliminary searches.

Conflicts of interest - None declared.

INPLASY registration number: INPLASY202460098

Amendments - This protocol was registered with the International Platform of Registered Systematic Review and Meta-Analysis Protocols (INPLASY) on 24 June 2024 and was last updated on 29 June 2024.

INTRODUCTION

Review question / Objective In patients with moderate-to-severe atopic dermatitis, which drug therapy is the most effective at reducing the risk of flare after 12 weeks of therapy compared to control?

Rationale A primary goal of atopic dermatitis treatment is to prevent and/or reduce the occurrence of flares. It is important to analyze the evidence to understand which therapies are the most effective at doing this.

Condition being studied Atopic dermatitis, also known as eczema, is a chronic inflammatory skin condition characterized by dry, itchy, and red skin. It is commonly found in infants and children but can persist into adulthood. The exact cause is unknown, but it is believed to involve a

combination of genetic, environmental, and immunological factors.

METHODS

Participant or population Moderate-to-severe atopic dermatitis patients.

Intervention Drug therapy (topicals and systemic agents).

Comparator Control (placebo).

Study designs to be included Double-blind randomized control trials.

Eligibility criteria Studies must have provided a Kaplan-Meier curve or cumulative risk curve.

Patients must be free of signs and symptoms of atopic dermatitis flare (stabilized), at the time of the therapy.

Primary treatment must be given in a standardized manner (e.g., at specified time intervals and not as needed).

Information sources Pubmed, CENTRAL, and Embase. Clinicaltrials.gov and WHO ICTRP can be searched for additional information.

Main outcome(s) Risk ratio of flare after 12 weeks of therapy.

Additional outcome(s) Relative risk reduction.

Quality assessment / Risk of bias analysis The Covidence systematic review platform will be used.

Strategy of data synthesis IPD will be reconstructed from Kaplan-Meier survival curves of time to flare or cumulative risk curves. IPD will then be analyzed using MetaInsight by doing at bayesian network meta-analysis of dichotomous outcomes.

Subgroup analysis None.

Sensitivity analysis Deviance, uncertainty, and reliability measures will be provided by MetaInsight and these will be analyzed.

Language restriction English.

Country(ies) involved United States of America.

Keywords atopic dermatitis; flare; relapse.

Contributions of each author

Author 1 - Dana Jaalouk.

Email: danajaalouk@hotmail.com

Author 2 - Jasmine Humeda.

Author 3 - Anika Pulumati.

Author 4 - Yanci A. Algarin.

Author 5 - Keyvan Nouri.