

INPLASY2024110127

doi: 10.37766/inplasy2024.11.0127

Received: 30 November 2024

Published: 30 November 2024

Bai, MY; Wang, HW; Cao, Y; An, SB.

Corresponding author:

Huawei Wang

aokayi@126.com

Author Affiliation:

Liaoning University Of Traditional Chinese Medicine.

ADMINISTRATIVE INFORMATION**Support** - None.**Review Stage at time of this submission** - Preliminary searches.**Conflicts of interest** - None declared.**INPLASY registration number:** INPLASY2024110127**Amendments** - This protocol was registered with the International Platform of Registered Systematic Review and Meta-Analysis Protocols (INPLASY) on 30 November 2024 and was last updated on 10 February 2024.**INTRODUCTION**

Review question / Objective To explore the insomnia detection rate and its influencing factors in Chinese tumour patients, provide reference for reducing the incidence of insomnia in Chinese tumour patients. The study types are observational studies such as cohort study, case-control study and cross-sectional study.

Condition being studied Insomnia is a common health problem among cancer patients, which is not only a physical problem but also a psychological problem. Insomnia is one of the most prominent complaints among patients with cancer. It is defined as difficulty falling asleep, trouble staying asleep, early awakening, or nonrestorative sleep. The prevalence of insomnia in patients with cancer varies from 30 to 60%, which is considerably higher than that in the general population. In addition, many cancer patients do not appear to report symptoms of insomnia, assuming them to be a minor problem

compared to a cancer diagnosis or treatment despite its prevalence and clinical significance, which results in insomnia remaining untreated. Insomnia and subsequent sleep disturbances can lead to fatigue, psychological disorders, and immunosuppression, which can significantly impair the quality of life and even affect the course of the disease.

METHODS

Participant or population The study population was Chinese tumour patients (≥ 18 years of age).

Intervention Exposures are Risk Factors for insomnia.

Comparator There will be no comparator as this is descriptive study.

Study designs to be included The included study designs are observational studies such as cohort study, case-control study and cross-sectional study.

Eligibility criteria

Inclusion Criteria:

- (1) Study population consists of Chinese cancer patients aged 18 years and older;
- (2) Study aims to investigate the incidence of insomnia and its influencing factors;
- (3) Outcome measures include the detection of insomnia using relevant sleep assessment tools such as the Pittsburgh Sleep Quality Index (PSQI) ;
- (4) Study type is cross-sectional;
- (5) Literature provides calculated or convertible odds ratios (OR) with 95% confidence intervals.

Exclusion Criteria:

- (1) Literature in languages other than Chinese or English;
- (2) Duplicate publications or literature with inaccessible raw data;
- (3) Literature unrelated to the research objectives;
- (4) Animal experiments, reviews, commentaries, or report-type literature;
- (5) Literature of poor quality.

Information sources PubMed,EmBase,Web of Science,Sinomed,Cochrane Library,CNKI,Wanfang Date and VIP.

Main outcome(s) Detection of insomnia in Chinese tumour patients.

Quality assessment / Risk of bias analysis The data were extracted and the literature quality was evaluated according to the cross-sectional study quality assessment criteria of Agency for Healthcare Research and Quality(AHRQ).

2 researchers used AHRQ to evaluate the included studies. In case of disagreement, a third researcher assisted in the judgement.The AHRQ has a total of 11 entries, and based on the total score the Studies were classified as low quality studies (0-3 points), medium quality studies (4-7 points) and high quality studies (8-11 points).

Strategy of data synthesis Statistical analysis was performed using Stata version 12.0 for Windows. Effect sizes were assessed using ORs of factors influencing insomnia in oncology patients and their 95% CIs to assess the correlation between specific factors and the occurrence of insomnia in oncology patients. Heterogeneity between studies was estimated using Cochrane's Q-test and I²for heterogeneity test, P \geq 0.1, I²< 50% suggests that there is no significant statistical heterogeneity, and a fixed-effects model was used; P<0.1, I² \geq 50% suggests that there is statistical heterogeneity a random-effects model was chosen for the combined analysis. Sensitivity analyses were performed by comparing the differences

between the fixed-effects model and the random-effects model combined. We evaluated the potential publication bias by funnel plots supplemented by the Egger regression asymmetry test. p<0.05 was considered statistically significant.

Subgroup analysis None.

Sensitivity analysis Sensitivity analyses were performed by comparing the differences between the fixed-effects model and the random-effects model combined.

Country(ies) involved China.

Keywords Insomnia; Tumour patients; Relevance ratio; Root cause analysis; Meta-analysis.

Contributions of each author

Author 1 - Mingyuan Bai.

Email: baimy0513@126.com

Author 2 - Huawei Wang.

Email: aokayi@126.com

Author 3 - Yun Cao.

Email: cy0219019@163.com

Author 4 - Sibao An.

Email: 1849408122@qq.com