

# INPLASY

INPLASY202480007

doi: 10.37766/inplasy2024.8.0007

Received: 01 August 2024

Published: 01 August 2024

**Corresponding author:**

Rebecca Pillai Riddell

rpr@yorku.ca

**Author Affiliation:**

Department of Psychology, Faculty of Health, York University, Toronto, Ontario, Canada; Department of Psychology, Neuroscience & Behaviour, McMaster University, Hamilton, Ontario, Canada.

## Patterns of heart rate variability (HRV) responses to acute skin-breaking procedures in neonates: A systematic review and meta-analysis

Bucsea, O.; Jasim, S.; Cohen, E.; Schmidt, L.; Pillai Riddell, R.

**ADMINISTRATIVE INFORMATION**

**Support** - Dr. Rebecca Pillai Riddell, a core faculty member of the Clinical-Developmental Psychology program at York University, is responsible for initiating and managing the review. This research is funded by operating grants from the Canadian Institutes of Health Research Natural Sciences and Engineering Research Council, and Social Sciences and Humanities Research Council to Dr. Pillai Riddell. Ms. Bucsea received support from the Canadian Institutes of Health Research (CIHR), Ontario Graduate Scholarship (OGS), and Canadian Psychological Association (CPA).

**Review Stage at time of this submission** - Formal screening of search results against eligibility criteria.

**Conflicts of interest** - None declared.

**INPLASY registration number:** INPLASY202480007

**Amendments** - This protocol was registered with the International Platform of Registered Systematic Review and Meta-Analysis Protocols (INPLASY) on 01 August 2024 and was last updated on 30 April 2025.

**INTRODUCTION**

**Review question / Objective** The overall objective of this systematic review is to synthesize the literature examining preterm and full-term newborns' acute heart rate variability (HRV) responses to acute skin-breaking procedures. It will address the following research question: What normative patterns of HRV responses do newborns exhibit during acute painful procedures?

**Rationale** Early neonatal exposure to repeated stressors, such as painful medical procedures in hospital settings, has been linked to impaired autonomic nervous system (ANS) stress responses. Furthermore, other clinical variables,

such as prematurity, have been associated with an underdeveloped ANS, increasing the vulnerability of hospitalized newborns to stress exposure. Heart rate variability (HRV) measures capture the balance of both branches of the ANS (sympathetic nervous system [SNS] and parasympathetic nervous system [PNS]), thus providing a non-invasive measure of stress reactivity and regulation. However, in newborns, heart rate has been more widely used to index cardiac stress responding to acute stress and pain in clinical settings. Although HRV measures are widely used to capture stress regulation in older ages, they need to be better understood in newborns and thus are currently being underused to measure acute stress physiology in healthcare settings. This systematic literature review seeks to examine patterns of HRV

responses to acute pain in newborns and variables that may influence these responses, such as gestational and postnatal age, as well as phase of the acute pain-related stress response (reactivity vs. regulation).

**Condition being studied** This review focuses on newborn infants exposed to any acute skin-breaking procedures in the first month of life. Acute HRV responses during these procedures capturing newborns' pain-related distress reactivity and regulation patterns is the main outcome of interest. The influence of age (i.e., gestational age and postnatal age) on the main outcome of interest will be examined.

## METHODS

**Search strategy** A search strategy was developed in collaboration with an academic librarian. The following databases were searched: Medline (PudMed), Embase, CINAHL, PsycINFO (Proquest and Ovid), Web of Science, and Scopus. The search was conducted on May 6, 2024 and updated on April 15, 2025 using the following search strategy:

newborn\* OR new-born\* OR neonate\* OR infant\* OR baby OR babies OR preterm OR pre-term OR prematurity OR premature OR "full term" OR "full-term" AND

pain\* OR nociceptive OR stress\* OR distress\* OR noxious\* AND

"heart rate variability" OR "heart-rate variability" OR HRV OR "respiratory sinus arrhythmia" OR RSA OR "vagal tone" OR "vagal withdrawal" OR "vagal regulation" OR "vagal reactivity" OR "NIPE" OR "Newborn Infant Parasympathetic Evaluation" OR "Newborn and Infant Parasympathetic Evaluation" OR "interbeat interval\*" OR "inter-beat interval\*" OR IBI OR "beat-to-beat interval\*".

**Participant or population** The review will focus on all preterm and full-term newborns in the first month of life. Newborns exposed to additional risk factors alongside prematurity (e.g., brain injury) will not be considered given the confounding influence of these variables on HRV activity.

**Intervention** All preterm and full-term newborn samples undergoing an acute skin-breaking procedures (e.g., heel lance, venipuncture, injection) will be included. For RCTs investigating the effects of an intervention on newborns' acute HRV responses which is outside the scope of the

current review, only the participants included in the control/standard care group will be considered.

**Comparator** Not applicable.

**Study designs to be included** Any original empirical article (both observational and RCT) will be considered eligible for inclusion. Review articles, commentaries, conference proceedings, case studies, and dissertations will not be included.

**Eligibility criteria** Studies that meet the following eligibility criteria are being selected for inclusion: 1) original empirical article; 2) study conducted in either preterm or full-term newborns within the first month of life; 3) includes an acute skin-breaking procedure that is considered painful to newborns; and 4) HRV measures are being collected before and during the acute procedure.

Studies are not excluded based on language, time frame, country or setting. Only peer-reviewed published studies will be included.

**Information sources** The following databases were searched: Medline (PudMed), Embase, CINAHL, PsycINFO (Proquest and Ovid), Web of Science, and Scopus. Google Scholar alerts are also being received weekly to identify additional eligible studies. Included and excluded abstracts from Google Scholar alerts will be tracked and included in the PRISMA flow chart. The reference lists of relevant reviews/chapters are also being manually inspected to ensure completeness.

**Main outcome(s)** The main outcomes of interest will include:

- Newborn HRV responses before, during, and after an acute skin-breaking procedure. Time-domain, frequency-domain, and non-linear types of HRV metrics will be included, as well as the Newborn and Infant Parasympathetic Evaluation (NIPE) index.

**Additional outcome(s)** N/A.

**Data management** Three authors (OB, SJ, EC) will be responsible for conducting title/abstract screening and full-text review to identify articles that meet criteria for inclusion. Disagreements will be discussed with senior author RPR. Data extraction will then be performed by the same three authors using a data extraction template developed in consultation with RPR. Inter-rater reliability will be assessed and tracked during the title/abstract screening, full-text review, and data extraction processes and disagreements will be

resolved via discussion and consultation with RPR, as needed.

The following data will be extracted from each study:

- Participant data, such as gestational age at birth, postnatal age at time of study, sex, average birth weight, illness severity, sample size, country of recruitment, setting of data collection, study beginning and end dates, inclusion/exclusion criteria, and details of the methodology and painful stimuli (e.g. type of painful procedure, description of procedure, time duration of procedure, description of baseline activity/handling, infant body positioning during study, pain management received during painful procedure).
- Duration of measurement for HRV measures (e.g. 15 seconds, 30 seconds, 1 minute)
- Information on newborn sample-specific HRV measures (i.e., how measures were recorded, edited, calculated, and interpreted)
- Descriptive statistics for newborn HRV measures (absolute values or changes from baseline)
- Results of statistical analyses examining trends in HRV responses during the acute painful procedure
- Any covariates and/or moderators (i.e., brain injury, prenatal exposure, birthweight, degree of prematurity, use of sucrose, medication status in inclusion/exclusion criteria, infant state during baseline [alert, drowsy, etc.], illness severity) examined in relation to the HRV responses
- Methodological limitations and future recommendations pertaining to HRV research in newborns

Covidence software and Microsoft Excel will be used to record inclusion/exclusion decisions throughout the systematic review process, as well as for data extraction and risk of bias judgements. RevMan software will be used for statistical analyses.

**Quality assessment / Risk of bias analysis** Three authors (OB, SJ, EC) will independently assess the risk of bias of individual included studies using an adapted risk of bias tool developed for the current review and informed by the NIH National Heart, Blood, and Lung Institute Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies, the Risk of Bias in Non-Randomized Studies – of Exposure (ROBINS-E) tool and published guidelines pertaining to best practices in the measurement and analysis of heart rate variability (HRV) outcome measures. Inter-rater reliability will be assessed, and disagreements will be resolved via discussion and consultation with the senior author (RPR). Assessments of risk of bias will inform overall data synthesis.

**Strategy of data synthesis** Each study's participant characteristics, acute procedure, study methodology, and study quality will be evaluated, as well as any outcomes and statistical analyses that pertain to the research question (i.e., patterns of newborn HRV responses to acute skin-breaking procedures). Given that several variables are expected to influence the pattern of HRV responses, findings will be stratified by 1) type of HRV measure; 2) acute pain phase (reactivity vs. regulation); and 3) gestational and postnatal age.

To address the first aim, standardized mean differences (Cohen's d) will be used to examine differences in HRV responses (continuous outcome) across the phases of acute responding within a procedure (baseline to reactivity, and baseline to regulation). These effect sizes will then be meta-analysed by calculating the weighted average using the Inverse Variance method. A 95% confidence interval using the Wald-type method will also be computed for the weighted average. A random-effects model will be performed. I<sup>2</sup> coefficients will be used to assess the heterogeneity among effect sizes. These statistical analyses will all be conducted in RevMan. When studies cannot be included in the quantitative analysis due to a lack of reporting of necessary statistical coefficients, they will be described qualitatively.

The most recent PRISMA guidelines (Page et al., 2021) will be followed when reporting the review findings.

**Subgroup analysis** Patterns of HRV responses to acutely painful procedures will be examined separately based on HRV outcomes reflecting the parasympathetic nervous system (PNS) activity with or without the combined influence of the sympathetic nervous system (SNS); acute pain response phase (reactivity and regulation); and gestational and postnatal age.

**Sensitivity analysis** Should considerable heterogeneity emerge across study effect sizes, studies with outlier effect sizes will be narratively inspected for sample (i.e., medication status, illness severity, infant state), methodological (i.e., type of acute painful procedure), or statistical (i.e., outcomes reported) differences.

**Language restriction** No language restrictions were imposed in the search strategy.

**Country(ies) involved** Canada.

---

**Keywords** Newborn, pain, acute, skin-breaking, heart rate variability, HRV.

**Dissemination plans** Review findings will be published in a peer-reviewed journal and presented at academic conferences.

#### **Contributions of each author**

Author 1 - Oana Bucsea – responsible for conceiving the review; designing the review; coordinating the review; data collection; data management; data analysis; data interpretation; writing the protocol or review.

Email: obucsea@yorku.ca

Author 2 - Sara Jasim – responsible for data collection; data management.

Email: sjasim@yorku.ca

Author 3 - Estreya Cohen - responsible for data collection; data management.

Email: estco@yorku.ca

Author 4 - Schmidt, L.

Author 5 - Rebecca Pillai Riddell – responsible for conceiving the review; designing the review; writing the protocol or review; data interpretation; providing funding.

Email: rpr@yorku.ca

#### **References**

Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ*. 2021 Mar 29;372.