

**Outcomes of revision surgery following failed total ankle arthroplasty: systematic review with narrative synthesis**

INPLASY202660048

doi: 10.37766/inplasy2026.6.0048

Received: 9 June 2026

Published: 10 June 2026

Kaiser, S; Kvarda, P.

**Corresponding author:**

Svenja Kaiser

svenja.kaiser@usb.ch

**Author Affiliation:**

University Hospital Basel.

**ADMINISTRATIVE INFORMATION****Support** - None.**Review Stage at time of this submission** - The review has not yet started.**Conflicts of interest** - None declared.**INPLASY registration number:** INPLASY202660048**Amendments** - This protocol was registered with the International Platform of Registered Systematic Review and Meta-Analysis Protocols (INPLASY) on 10 June 2026 and was last updated on 10 June 2026.**INTRODUCTION**

**R** **Review question / Objective** n adult patients who have undergone primary total ankle arthroplasty (TAA) and subsequently developed implant failure, what are the clinical outcomes — including implant survival, re-revision rates, functional outcomes, and complication profiles — of revision surgical procedures (revision TAA, revision ankle arthrodesis, revision tibiototalcalcaneal fusion)?

**P (Population):** Adult patients ( $\geq 18$  years) with failed primary TAA requiring revision surgery

**I (Intervention):** Any revision surgical procedure (RTAA, RAA, RTTC, or other salvage procedure)

**C (Comparison):** Comparison between revision modalities where available; single-arm studies also eligible

**O (Outcomes):** Primary: implant survival, re-revision rate, conversion to arthrodesis/amputation, failure time and cause

Secondary: PROMs (EFAS, AOFAS, VAS), complication rates, radiographic outcomes

**S (Study design):** RCTs, prospective/retrospective cohorts, registry studies, case series.

**Rationale** With the increasing number of primary TAA procedures, the absolute number of revision surgeries is also rising, and further increases are anticipated. Despite the increasing importance of TAA revisions, available data remains limited.

**Condition being studied** End-stage ankle osteoarthritis results in severe pain, significant functional limitations, and diminished quality of life. When conservative management is unsuccessful, surgical intervention becomes necessary. Ankle arthrodesis (AA) has traditionally been considered the gold standard, providing reliable pain relief and

durable outcomes but changes of biomechanical function. Total ankle arthroplasty (TAA) offers a motion-preserving alternative.

## METHODS

**Search strategy** Databases: MEDLINE (via PubMed), Embase (via Ovid), Cochrane Central Register of Controlled Trials (CENTRAL)

Additional sources: Reference lists of included studies (backward citation tracking); forward citation tracking via Web of Science; ClinicalTrials.gov; manual search of Foot & Ankle International, Bone & Joint Journal, Foot and Ankle Surgery

PubMed search string:  
 ("total ankle arthroplasty"[MeSH] OR "total ankle replacement"[tiab] OR "ankle arthroplasty"[tiab] OR "ankle prosthesis"[tiab])  
 AND ("revision"[tiab] OR "re-revision"[tiab] OR "salvage"[tiab] OR "failed"[tiab] OR "failure"[tiab] OR "reoperation"[tiab])  
 AND ("survival"[tiab] OR "outcome"[tiab] OR "complication"[tiab] OR "arthrodesis"[tiab] OR "fusion"[tiab] OR "periprosthetic"[tiab]).

**Participant or population** Adult patients ( $\geq 18$  years) who underwent revision surgery after failed primary TAA.

**Intervention** Surgical interventions: total ankle arthroplasty, ankle arthrodesis; tibiototalcalcaneal fusion.

**Comparator** Where available, comparison between different revision modalities: total ankle arthroplasty, ankle arthrodesis; tibiototalcalcaneal fusion. Single arm studies without a comparator group are also eligible for inclusion.

**Study designs to be included** Randomised controlled trials, prospective and retrospective cohort studies, registry-based studies, case series with a minimum of 5 revision surgeries. conference abstracts, letters to the editor, editorials and narrative reviews will be excluded.

**Eligibility criteria** Inclusion: Adults ( $\geq 18$  years) with failed primary Total ankle arthroplasty (TAA) requiring revision surgery. Minimum 5 revision surgeries. At least one reported outcome of interest. Full text available. Exclusion: primary TAA studies without revision cohort, isolated polypropylene exchange, less than 5 surgeries, abstracts without full text, duplicate datasets.

**Information sources** Electronic databases: MEDLINE (PubMed), Embase (Ovid), Cochrane CENTRAL. Additional sources: Backward and forward citation tracking of included studies, manual search of Foot & Ankle International, Bone & Joint Journal.

**Main outcome(s)** Primary:

1. Implant survival (Kaplan-Meier estimates at 1, 3, 5, and 10 years where available)
2. Re-revision rate (further revision surgery of any kind)
3. Conversion to arthrodesis or amputation
4. Time to failure, causes of failure.

**Additional outcome(s)** Secondary:

1. Patient-reported outcome measures: EFAS score, AOFAS Ankle-Hindfoot Score, VAS pain score
2. Complication rates (wound complications, deep infection, periprosthetic fracture, nerve injury)
3. Radiographic outcomes (alignment, osteolysis, subsidence)
4. Risk factors for failure (patient-level and implant-level).

**Data management** Search results from all databases will be exported and imported into Zotero for reference management. Duplicates will be removed. Study selection will be performed using Rayyan. Standardised data extraction forms (Microsoft Excel) will be used to collect data from included studies. All extracted data will be stored securely and made available upon reasonable request.

**Quality assessment / Risk of bias analysis** The methodological quality of all included non-randomised studies will be assessed using the Methodological Index for non-randomised studies (MINORS).

**Strategy of data synthesis** Due to anticipated clinical and methodological heterogeneity across included studies (differences in study design, patient population, follow-up duration, outcome definitions), a formal meta-analysis will not be performed. Data will be synthesised narratively and presented in structured evidence tables.

**Subgroup analysis** Subgroup analyses are planned for: (1) revision indication (septic vs. aseptic); (2) implant system used at revision; (3) follow-up duration (short-term 5 years). These analyses are contingent on sufficient data being available in the included studies.

---

**Sensitivity analysis** A sensitivity analysis will be performed by restricting the analysis to studies with a minimum follow-up of 24 months, to assess whether short-term studies disproportionately influence the overall findings. Additionally a sensitivity analysis will be conducted excluding studies rated as low methodological quality (MINORS Score less than 10/16 for non-comparative and 16/24 for comparative studies), to evaluate the robustness of results when restricted to higher-quality evidence only.

**Language restriction** English.

**Country(ies) involved** Switzerland.

**Other relevant information** This review is conducted as a systematic review with narrative synthesis. A formal meta-analysis was not planned due to anticipated heterogeneity in study design, patient population, implant systems and outcome reporting. The review follows the PRISMA 2020 guidelines. No ethical approval is required as this review is based solely on published literature. Any amendments to this protocol after registration will be documented transparently in the final publication.

**Keywords** total ankle arthroplasty; total ankle replacement; revision arthroplasty; ankle arthrodesis; tibiototalcalcaneal fusion; implant survival; salvage procedure; periprosthetic joint infection; aseptic.

**Dissemination plans** The results of this systematic review with narrative synthesis will be submitted for publication in a peer-reviewed orthopaedic journal (Minerva Medica).

#### **Contributions of each author**

Author 1 - Svenja Kaiser - Developed the protocol, performed title/abstract and full-text screening, conducted data extraction and quality assessment, drafted manuscript.

Email: svenja.kaiser@usb.ch

Author 2 - Peter Kvarda - Independently performed title/abstract and full-text screening, conducted data extraction and quality assessment, critically revised manuscript for important intellectual content.